

# METHODS FOR PREDICTING SERIOUS EYE DAMAGE AND IRRITATION

METHOD	PRINCIPLE OF THE TEST	APPLICABILITY DOMAIN	GHS CATEGORISATION
<b>DEFINED APPROACHES</b>			
<b>OECD Test Guideline 467: Defined Approaches for Serious Eye Damage and Eye Irritation, comprising two defined approaches for liquids (DAL1 and DAL2) and a defined approach for solids (DAS)</b>	<p>DAL1 combines information from three sources: (1) physicochemical properties of the test substance, (2) results from testing in OECD TG 437 BCOP assay using the laser light based opacitometer (LLBO), and (3) results from testing in OECD TG 492 (either the EpiOcular Eye Irritation Test or the SkinEthic Human Corneal Epithelium Eye Irritation Test). DAL2 combines information from two sources: (1) results from testing in OECD TG 437 BCOP assay using the LLBO and (2) OECD TG 491 Short Time Exposure assay.</p> <p>DAS combines information from two <i>in vitro</i> sources: (1) results from testing in OECD TG 437 BCOP assay using LLBO and (2) results from testing in OECD TG 492 (using the SkinEthic™ Human Corneal Epithelium Eye Irritation Test).</p>	DAL1 is applicable to neat non-surfactant liquids, DAL2 is applicable to neat and diluted non-surfactant liquids, and DAS is applicable to neat solids. For all approaches, the applicability domain of each test guideline used within the defined approach should also be considered.	The approaches can be used for the identification of substances causing serious eye damage (GHS Cat 1), causing eye irritation (GHS Cat 2), and not requiring classification for eye irritation or serious eye damage (GHS No Cat). Each approach can be used as a full replacement for the Draize rabbit eye irritation test.
<b>RECONSTRUCTED THREE-DIMENSIONAL HUMAN TISSUE ASSAYS</b>			
<b>OECD Test Guideline 492: Reconstructed Human Cornea-like Epithelium Test Method for Identifying Chemicals Not Requiring Classification and Labelling for Eye Irritation or Serious Eye Damage (e.g. EpiOcular™, SkinEthic™, LabCyte, and MCTT HCE™)</b>	The test substance is applied to reconstructed tissue from human cells, which have been cultured to form a stratified, highly differentiated squamous epithelium that is morphologically similar to that found in the human cornea. Cell viability (MTT or WST-8 assay) is used to predict toxicity.	This assay is applicable to substances and mixtures and to solids, liquids, semi-solids, and waxes.	This assay can be used for the identification of substances not requiring classification for eye irritation or serious eye damage (GHS No Cat).
<b>OECD Test Guideline 492B: Reconstructed Human Cornea-like Epithelium Test Method for Eye Hazard Identification (SkinEthic™)</b>	The test substance is applied to reconstructed tissue from human cells (as in OECD TG 492). Depending on whether the test substance is a solid or a liquid, cell viability is assessed at two or three exposure times, respectively.	This assay is applicable to substances and mixtures and to solids, liquids, semi-solids, and waxes.	This assay can be used for the identification of substances causing serious eye damage (GHS Cat 1), causing eye irritation (GHS Cat 2), and not requiring classification for eye irritation or serious eye damage (GHS No Cat). This method can be used as a full replacement for the Draize rabbit eye irritation test.
<b>OECD Test Guideline 494: Vitrigel-Eye Irritancy Test Method for Identifying Chemicals Not Requiring Classification and Labelling for Eye Irritation or Serious Eye Damage</b>	Human corneal epithelium models fabricated in a collagen vitrigel membrane are exposed to a test substance. Damage to the barrier function of the models is assessed by analysing time-dependent changes in transepithelial electrical resistance values.	This assay is applicable to substances and mixtures with a pH > 5 (based on 2.5% weight/volume preparation). It is not applicable to solids.	This assay can be used for the identification of substances not requiring classification for eye irritation or serious eye damage (GHS No Cat).
<b>ORGANOTYPIC EX VIVO ASSAYS</b>			
<b>OECD Test Guideline 437: Bovine Corneal Opacity and Permeability (BCOP) Test Method for Identifying i) Chemicals Inducing Serious Eye Damage and ii) Chemicals Not Requiring Classification for Eye Irritation or Serious Eye Damage</b>	The test substance is directly applied to cows' eyes obtained as by-products from abattoirs. Corneal opacity (measured quantitatively as the amount of light transmission through the cornea) and permeability (measured quantitatively as the amount of sodium fluorescein dye that passes across the full thickness of the cornea) are measured. Optional histopathology can be conducted for additional information.	This assay is applicable to solids, liquids (including semi-solids, creams, and waxes), and mixtures.	This assay can be used for the identification of substances causing serious eye damage (GHS Cat 1) and substances not requiring classification for eye irritation or serious eye damage (GHS No Cat).
<b>OECD Test Guideline 438: Isolated Chicken Eye (ICE) Test Method for Identifying i) Chemicals Inducing Serious Eye Damage and ii) Chemicals Not Requiring Classification for Eye Irritation or Serious Eye Damage</b>	The test substance is directly applied to chickens' eyes obtained as by-products from abattoirs. Corneal swelling, opacity, and fluorescein retention are assessed.	This assay is applicable to solids (which may be soluble or insoluble in water), liquids, emulsions, and gels.	This assay can be used for the identification of substances causing serious eye damage (GHS Cat 1) and substances not requiring classification for eye irritation or serious eye damage (GHS No Cat).
<b>CYTOTOXICITY AND CELL FUNCTION BASED IN VITRO ASSAYS</b>			
<b>OECD Test Guideline 460: Fluorescein Leakage (FL) Test Method for Identifying Ocular Corrosives and Severe Irritants</b>	Epithelial monolayer Madin-Darby canine kidney cells are cultured on permeable inserts. The test chemical is applied for 1 minute and then removed. Next, the non-toxic, highly fluorescent sodium fluorescein dye is added, and the amount of dye that passes through the cell layer is measured spectrophotometrically and used to predict toxicity.	This assay is applicable to water-soluble chemicals or mixtures. There are limitations for coloured or highly viscous substances. (However, predictivity is improved by increasing the number of wash steps.) It is not applicable to strong acids and bases, cell fixatives, or highly volatile substances.	This assay can be used for the identification of substances causing serious eye damage (GHS Cat 1).

METHOD	PRINCIPLE OF THE TEST	APPLICABILITY DOMAIN	GHS CATEGORISATION
<b>OECD Test Guideline 491: Short Time Exposure (STE) /<i>In Vitro</i> Test Method for Identifying i) Chemicals Inducing Serious Eye Damage and ii) Chemicals Not Requiring Classification for Eye Irritation or Serious Eye Damage</b>	This test measures cell viability (via the MTT assay) of Statens Seruminstitut Rabbit Cornea (SIRC) corneal epithelial cells in 96 well plates. As compounds are generally cleared from human eyes in 1 to 2 minutes and from rabbit eyes in 3 to 4 minutes, this test requires a 5-minute exposure.	This assay is applicable to test chemicals that are soluble in saline, DMSO, or mineral oil.	This assay can be used for the identification of substances causing serious eye damage (GHS Cat 1) and substances not requiring classification for eye irritation or serious eye damage (GHS No Cat).
<b>MACROMOLECULAR MATRIX ASSAYS</b>			
<b>OECD Test Guideline 496: <i>In Vitro</i> Macromolecular Test Method for Identifying Chemicals Inducing Serious Eye Damage and Chemicals Not Requiring Classification for Eye Irritation or Serious Eye Damage</b>	The test substance is directly applied to an <i>in chemico</i> macromolecular matrix model composed of lipids, proteins, glycoproteins, carbohydrates, and low molecular weight substances that model the cellular structure of the human corneal epithelium. An increase in optical density is used to predict the ocular hazard effects of chemicals.	This assay is applicable to solids (which may be soluble or insoluble in water) and liquids (which may be viscous or non-viscous) whose 10% solution/dispersion has a pH in the range $4 \leq \text{pH} \leq 9$ . There are some limitations for intensely coloured chemicals, chemicals that cause salting-out precipitation, high concentrations of some surfactants, and highly volatile chemicals. It is also applicable to mixtures.	This assay can be used for the identification of substances causing serious eye damage (GHS Cat 1) and/or substances not requiring classification for eye irritation or serious eye damage (GHS No Cat).

For more information on these methods, please see our publication, Clippinger AJ et al. Human-relevant approaches to assess eye corrosion/irritation potential of agrochemical formulations. *Cutan Ocul Toxicol.* 2021;40(2):145-167.

For more information on tiered testing strategies for serious eye damage and eye irritation, please see the following publications:

- Organisation for Economic Co-operation and Development. 2019. Guidance document on integrated approaches to testing and assessment (IATA) for serious eye damage and eye irritation. No 263. Series on Testing and Assessment.
- US Environmental Protection Agency. 2015. Use of an alternate testing framework for classification of eye irritation potential of EPA pesticide products.
- European Chemicals Agency. 2017. Guidance on information requirements and chemical safety assessment. Chapter R.7a: Endpoint specific guidance. Version 6.0. See R.7.2.8–R.7.2.11.